



© Igor Mojzes | Dreamstime.com

Is Beauty Truly In The Eye Of The Beholder? Evidence-Based Ideal Smile Esthetics

A peer-reviewed article written by Maria L. Geisinger, DDS, MS, and Hussein Basma, DDS, DESS, MS

PUBLICATION DATE:	MAY 2021
EXPIRATION DATE:	APRIL 2024

EARN

3 CE
CREDITS

Is beauty truly in the eye of the beholder? Evidence-based ideal smile esthetics

ABSTRACT

A smile is a universal greeting and translates into all languages. Smiling individuals are judged as more pleasant and trustworthy, and the act of smiling actually releases endorphins that improve the mood of the person who smiles. As dentists, we are trained to create, maintain, and protect the ideal smile, but what should our goals be in achieving that “gold standard” of beauty?

The essential components of an ideal smile involve the relationship between three primary components: the teeth, the lip framework, and the gingival scaffold. While beauty may be in the eye of the beholder, factors that allow assessment of overall smile esthetics include tooth width/height ratio, shape, position, quality of restoration, and general arrangement of the dentition, especially of the anterior teeth, upper lip position, buccal corridor, visibility of teeth, and amount of gingival display. These factors are considered in concert and usually judged esthetically as a group. It is considered that in the composition of a beautiful smile, the form balance, symmetry, and relationship of these elements make it attractive or unattractive. This course seeks to present the current data regarding ideal smile components and the differences regarding ideal smile components based upon age, gender, and race/ethnicity to allow dentists to create personalized road maps and help their patients achieve vibrant smiles.

EDUCATIONAL OBJECTIVES

At the conclusion of this course, the reader should be able to:

1. List the components of a smile and discuss their roles in optimal smile esthetics
2. Describe the step-by-step approach to evaluating a patient's smile, including assessing tooth shape/shade, gingival display, and lip length/mobility
3. Understand the role of patient-based characteristics on acceptable esthetics and ideal smile components
4. Discuss personalized treatment options to achieve ideal smile esthetics based upon underlying diagnoses in each patient



Dental Academy of Continuing Education™

Go online to take this course.
DentalAcademyofCE.com

QUICK ACCESS CODE 21002

This continuing education (CE) activity was developed by Endeavor Business Media with no commercial support.

This course was written for dentists, dental hygienists, and dental assistants, from novice to skilled.

Educational methods: This course is a self-instructional journal and web activity.

Provider disclosure: Endeavor Business Media neither has a leadership position nor a commercial interest in any products or services discussed or shared in this educational activity. No manufacturer or third party had any input in the development of the course content.

Requirements for successful completion: To obtain three (3) CE credits for this educational activity, you must pay the required fee, review the material, complete the course evaluation, and obtain an exam score of 70% or higher.

CE planner disclosure: Laura Winfield, Endeavor Business Media dental group CE coordinator, neither has a leadership nor commercial interest with the products or services discussed in this educational activity. Ms. Winfield can be reached at lwinfield@endeavorb2b.com.

Educational disclaimer: Completing a single continuing education course does not provide enough information to result in the participant being an expert in the field related to the course topic. It is a combination of many educational courses and clinical experience that allows the participant to develop skills and expertise.

Image authenticity statement: The images in this educational activity have not been altered.

Scientific integrity statement: Information shared in this CE course is developed from clinical research and represents the most current information available from evidence-based dentistry.

Known benefits and limitations of the data: The information presented in this educational activity is derived from the data and information contained in the reference section.

Registration: The cost of this CE course is \$59 for three (3) CE credits.

Cancellation and refund policy: Any participant who is not 100% satisfied with this course can request a full refund by contacting Endeavor Business Media in writing.

Provider information:

Dental Board of California: Provider RP5933. Course registration number CA code: 03-5933-21002. Expires 7/31/2022. “This course meets the Dental Board of California’s requirements for three (3) units of continuing education.”



Endeavor Business Media is a nationally approved PACE program provider for FAGD/MAGD credit.

Approval does not imply acceptance by any regulatory authority or AGD endorsement. 11/1/2019 to 10/31/2022.

Provider ID# 320452

AGD code: 780



Endeavor Business Media is designated as an approved Provider by the American Academy of Dental Hygiene, Inc. #AADHPNW (January 1, 2021-December 31, 2022). Approval does not imply acceptance by a state or provincial Board of Dentistry. Licensee should maintain this document in the event of an audit.

ADA CERP® Continuing Education Recognition Program

Endeavor Business Media is an ADA CERP-recognized provider.

ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of dental continuing education. ADA CERP does not approve or endorse individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry.

Concerns or complaints about a CE provider may be directed to the provider or to ADA CERP at ada.org/goto/cerp.



INTRODUCTION

Psychologists and scientists agree that the simple act of smiling can positively transform a person's life, as well as the lives of others. Hence, the smile can be considered one of the most fundamental factors to one's overall happiness and perceived beauty. The smile has become elevated to the forefront of the dental field with the increasing exposure by visual and entertainment media esthetic standards, decline in dental caries, and overall patient demand to improve their smiles. A study commissioned by the American Association of Orthodontists suggests that more than one third of US adults are unhappy with their smile esthetics.¹ This report also reviewed the social impact of decreased perceived esthetics, suggesting that 36% of those unhappy with their smiles believe they would have better social lives if they had better teeth.¹ Younger adults were particularly impacted; 48% of Americans ages 18–24 have untagged pictures of themselves on social media because they didn't like their smiles.¹

A smile involves the relationship between the three primary components: teeth, lip framework, and the gingival scaffold.² While beauty may be in the eye of the beholder, certain characteristics of each of these components are associated with perceived esthetics by dental professionals and laypeople. These characteristics include: tooth width/height ratio; tooth shape and position; quality of restorations; occlusal scheme and crowding; upper lip position; buccal corridor; visibility of teeth; and amount of gingival display.^{3–5} When assessing esthetics, dental health-care providers should consider such individual components in concert. Smiles are usually judged as a whole, and harmony of all components factor into the overall esthetics.^{5,6} It is considered that in the composition of a beautiful smile, the form, balance, symmetry, and relationship of these elements make it attractive or unattractive. This course will assess the individual components that contribute to perceived smile esthetics and will also present the ways in which the interaction of those components in a dynamic smile work to create ideal esthetics.




FIGURE 1: Common causes of compromised smile esthetics

WHAT MAKES A SMILE ESTHETIC? ORAL COMPONENTS OF AN IDEAL SMILE

Overall smile esthetics involve three oral tissues (teeth, lips, and gingivae) and their relationship with one another.² When assessing individuals who stated that they did not find their smiles ideally esthetic, complaints included all of the established components of a smile. Studies have demonstrated that tooth color was the most common smile component causing dissatisfaction among the subjects (27.9%), followed by tooth size (19.2%), tooth position (16.2%), tooth shape (15.0%), and lip shape (11.2%).⁷

Because of the broad range of complaints associated with compromised esthetics, an interdisciplinary approach to achieving ideal smile design allows for optimization of results. Employing an armamentarium of restorative techniques, orthodontic tooth movement, periodontal plastic surgery procedures, and facial esthetic procedures may be critical to address a combinatorial defect in smile esthetics. It is critical, therefore, that dental health-care professionals understand the individual components of a smile that are judged to be esthetic and mechanisms to maximize esthetics to improve their patients' smiles (figure 1).

TEETH Shade

Color is usually described according to the Munsell color space in terms of hue, value, and chroma. Hue is the attribute

of a color that enables the clinician to distinguish between different families of color, whereas value indicates the lightness of a color. Chroma is the degree of color saturation. In dentistry, the hue is characterized by a letter "A" through "D." The average shade for an untreated tooth is A3 and the lightest naturally occurring shade is generally considered to be B1.⁸ Tooth shade generally darkens as patients age^{9,10} and women are more likely to present with teeth of a lighter shade than men.¹⁰ It is generally accepted that patients prefer a lighter tooth shade, but this may be influenced by skin and lip color, age, and gender.^{11,12} Furthermore, other components of light reflection—such as translucency, fluorescence, and opalescence—may also influence the perceived esthetics.

Tooth size

Both tooth width:length and the width-to-length ratio of teeth are critical to achieving smile esthetics. The average vertical height of maxillary central incisors is 10.6 mm in males and 9.8 mm in females.^{13,14} Short clinical tooth crowns may be due to fracture, attrition, altered passive eruption, or gingival overgrowth. Extended clinical crowns may be due to gingival recession or improperly designed dental restorations. Similarly, the width:length ratio of teeth is critical for perceived esthetics, with the mean width:length ratio being 78% and a range of 75%–80% judged as esthetically acceptable.¹⁵

Anterior tooth inclination

Proclined maxillary incisors tend to reduce the incisor display, whereas uprighted or retroclined maxillary incisors generally increase the incisor display. Therefore, tooth inclination may contribute to inadequate or excessive tooth display.¹⁶

Occlusal plane orientation

The cant of the occlusal plane can affect the relationship between the maxillary teeth and the lower lip both at rest and during smile display.¹⁶ For example, a cant of the maxillary occlusal plane directed upward and anteriorly will result in maxillary incisal edges that do not follow the curvature of the lower lip, reducing esthetics in a smile.

Dental midline

The dental midline is an important focal point in an esthetic smile.¹⁷ Generally, a dental midline that corresponds with the facial midline—a line from the nasion to the base of the philtrum, or “cupid’s bow”—is deemed most esthetic, although significant deviation must be present for a smile to be judged as unattractive by dentists and laypeople.¹⁸ Therefore, mild midline discrepancies may be acceptable in cases with other esthetic smile components and a vertically oriented interproximal contact between the maxillary central incisors.¹⁸

Buccal corridor/transverse arch dimension

The buccal corridor is generally considered to be the negative or empty space between the buccal surfaces of the posterior teeth and the commissures of the mouth when smiling.^{19,20} The esthetic value of this space is equivocal with some data suggesting that dentures that completely fill the space are considered less esthetic and less natural in appearance and other literature suggesting that postorthodontic therapy that fills this space and results in a first molar to first molar smile is optimal.²⁰⁻²² A lack of buccal corridor fill also has not been determined to be detectable by laypersons as influencing esthetics.²² Transverse arch dimension affects the buccal corridor fill; a broad arch is more likely to fill the

buccal corridors than a narrow and constricted arch. Buccal corridor fill may also be influenced by the relative anteroposterior position of the maxilla in relation to the frame of the lips.^{16,19}

LIPS

Lip length

Lip length is generally measured in repose from the subnasale to the most inferior visible portion of the upper lip at the midline. Mean lip length is 23 mm in males and 20 mm in females and has been shown to increase with age.¹¹

Lip length also should be approximately equal in length with commissure height, i.e., the vertical distance between the commissures and a horizontal line from the subnasale point. During a full smile, the commissures of the mouth move outward and upward, and this creates the arched lip form considered esthetic. A short upper lip can result in lip incompetence, excessive gingival display, and/or a reverse resting upper lip line.²³

Lip mobility

In a full smile, the upper lip elevates to display the anterior teeth. On average, this lip mobility accounts for 80% of the original lip length, and maxillary incisal display at full smile is approximately 10 mm.²² While there is considerable interindividual variation in lip mobility, with a physiologic range of 2–12 mm,²⁴ women tend to have more mobility than men.²² Excessive lip mobility could result in excessive gingival display at full smile in cases with normal lip length.

Lip curvature

Lip curvature is characterized by the position of the upper lip at the midline relative to the position of the commissure in full smile. It may be upward, straight, or downward.^{21,25,26} Both laypersons and dentists judge upward and straight lip curvatures to be more esthetic than downward lip curvatures.²¹ Because lip curvature is determined by facial musculature, dental treatment alone may be limited in treating an individual with a downward lip in order to achieve an optimal smile.

Lip volume

Voluminous lips—generally resulting in greater maxillary incisor exposure at smiling, at rest, or while speaking—are considered critical to a current standard of esthetics.²⁷ Lip volume is influenced by anteroposterior positioning of teeth as well as anatomical variations in lip size.²⁸ Lip volume has been shown to decrease with age.²⁸ It is also important to note that the relative volume of the lower lip is generally greater than the upper lip, but upper lip augmentation procedures with surgical interventions and soft tissue fillers increased 60% and 312%, respectively, between 2000 and 2017.²⁹

GINGIVAL SCAFFOLD

Gingival color/texture

Inflammation, leading to gingival edema, can cause alterations in color, texture, and contour of the gingiva.³⁰ Healthy gingiva is generally considered to be coral pink in color and may have a stippled (or “tufted”) appearance. Gingival erythema and/or rolled gingival margins can be perceived as unesthetic.³¹

Papillary fill

Gingival tissues in the interdental areas, papillae, are considered most esthetic when they just fill the interdental space. Papillary overfill, as in the case of significant gingival inflammation, and missing or blunted gingival papillae are both considered unesthetic.³¹ Incomplete papillary fill, i.e., a “black triangle,” can be caused by interproximal bone loss, triangular tooth form, root divergence, open dental contacts, or as a result of resective periodontal surgery. If the distance from the interdental contact point to the crest of the interproximal bone is 5 mm, papillae are present 100% of the time, but the prevalence of papillae when the distance between the contact point and the interproximal crest is 6 mm is only 56%.³²

Gingival contours

The height of the gingival margins of anterior teeth and their relationship to one another is a component of esthetics. The height of gingival contour of the central incisors generally corresponds to those of the canines with the gingival margins of the

lateral incisors approximately 1–1.5 mm more coronal than those of the canines/central incisors. Discrepancies in these gingival margins may be caused by incisal attrition, ankylosis, altered passive eruption, dental malocclusion, or gingival recession.³³⁻³⁵

Gingival display

Upper lip position at full smile that is concordant with the gingival zeniths of the central incisors results in no gingival display.^{26,36,37} Excessive gingival display, sometimes called a “gummy smile,” may be categorized into one of five classes, ranging from class 1, where the lip is positioned well above the cervical portion of the maxillary teeth, to class 5, where the lip is positioned to provide full coverage of the maxillary teeth. While the ideal amount of gingival display is not uniform among populations, some investigators have shown that ideal lip position occurs when the lower margin of the upper lip aligns evenly with the gingival margin of the maxillary central incisors.^{26,36,37} However, other studies have demonstrated a combination of maxillary incisor visibility and gingival display up to 1 mm is judged equally or more attractive.^{38,39} The amount of gingival display considered acceptable may vary based upon subject age and gender and the societal norms within individual communities.^{18,40} Excessive gingival display is determined to be more detrimental to esthetics in men when compared to women.^{18,40} For instance, studies in some communities show that patients with gingival display in excess of 3 mm can be considered to have acceptable esthetics.^{18,40}

SMILE COMPONENT INTERACTION AND SYMMETRY

Overall facial attractiveness is driven by many factors, including the interaction of the individual component of smile esthetics, symmetry, and cultural/gender influences on perception.^{41,42} Facial attractiveness, including smile esthetics, drives the quality of dating partners, and attractive individuals are perceived to be more intelligent and nicer.⁴³⁻⁴⁶ However, quantification of facial attractiveness and smile esthetics as well as calibration of a single

standard for esthetics has proved elusive. Standards for these ideals also diverge between professionals and laypersons.⁴⁷ Overall, higher levels of facial and smile symmetry and meeting associated gender norms have been correlated to increased levels of attractiveness.^{48,49}

Smile symmetry

Smile symmetry, including the symmetry of individual smile components, e.g., tooth size and gingival contours, and the dynamic movement of the lips during smiling are key components of smile esthetics.⁵⁰ The presence of a unilateral lateral incisor width change of >1 mm was considered unesthetic by both dentists and laypersons, but bilateral changes of the same amount were not judged as harshly.⁵¹ Furthermore, large left-to-right differences in the relative positioning of the corners of the mouth in the vertical plane at full smile, such as is seen in patients with muscular tone impairment, has a significant negative impact on smile esthetics.^{52,53} Additionally, in these cases, an oblique commissural line, i.e., one that is off-parallel with the interpupillary line, may give the illusion of skeletal asymmetry and/or a transverse cant of the maxilla and negatively affect esthetics.¹⁶

Smile arc

Overall, the alignment of the maxillary incisal edges with the inner contour of the lower lip at full smile is considered to be ideal and is often described as “consonant.”⁵⁴ The curvature of the incisal edges and lip curvature tend to be more pronounced in younger smiles and flatten with age,⁵⁵ leading to the perceived desirability of such a relationship for optimal esthetics. Smile arcs were found to be flatter in orthodontically treated patients than in an untreated group with normal occlusions, which may inadvertently compromise overall esthetics.^{17,26,56}

ASSESSING SMILE ESTHETICS: CLINICAL AND PHOTOGRAPHIC EXAMINATION

When treating a patient for esthetic concerns, it is vital for the dental practitioner to understand what defines an esthetic smile. Is it truly in the eye of the beholder,

the dental health-care professional’s perception, or a combination of the two? In order to critically assess the individual smile components and their interaction, careful examination and documentation are necessary in both static and dynamic function. Full facial photographs at rest and full smile in a 1:1 image ratio can allow for measurements of dental midline alignment, lip length, lip mobility, gingival display, buccal corridor fill, smile cant, and consonant/nonconsonant smile arc. Intraoral measurements of incisor length, width:length ratios, gingival recession and/or overgrowth, and papillary fill should also be recorded during a comprehensive intraoral examination. Lastly, utilization of videography to assess dynamic components of smile esthetics may be helpful in patients with high esthetic concerns.

Emerging technologies, including machine learning and three-dimensional stereophotogrammetry, have been suggested to improve assessments and eliminate perception bias and manipulation.⁵⁷⁻⁵⁹ Currently, these technologies may not be reliable enough to fully assess facial attractiveness and smile esthetics. In their current form, they may also be very time consuming and technique sensitive, but they have been incorporated into research and may prove to be valuable tools as their use increases to include larger populations.

SMILES FROM MONA LISA TO JULIA ROBERTS: WHAT INFLUENCES VARYING STANDARDS OF SMILE ESTHETICS?

While we often consider certain characteristics to be universally appealing, facial attractiveness and smile esthetics are influenced by the individuals judging the ultimate result. It is well established that dental health-care professionals and plastic surgeons have a higher standard for facial esthetics than laypersons with similar demographics and socioeconomic status.^{60,61} Furthermore, age, gender, geographic divergence, and cultural norms may impact perception of smile esthetics, with men and older individuals being more tolerant of certain deviations from esthetic standards.⁶²⁻⁶⁷ For patients, their individual standards and self-perception affect their



FIGURE 2: Excessive gingival display (“gummy smile”) causes and interventions: esthetic crown lengthening and lip repositioning surgeries to address altered passive eruption and hypermobile upper lip, respectively.

preferences with regard to the importance of smile esthetics more so than the opinions of their dental health-care providers.⁶² Lastly, esthetic ideals are not static and changing parameters of ideal smile esthetics can be influenced by mass media, including TV, films, magazine, fashion, ads, and social media.^{68,69}

Given the myriad influences on perception of attractiveness and the relative importance of patients’ self-evaluation of their smile esthetics, it is important to thoroughly probe patients’ chief complaints when they present with esthetic concerns. The esoteric nature of “ideal” attractiveness also increases the importance of initial trials with reversible interventions, including digital modeling, to assess patient satisfaction with likely outcomes of therapy and to manage patient expectations for predictable results.

SMILES BY DESIGN: INTERVENTIONS TO ACHIEVE OPTIMAL SMILE ESTHETICS

Generally, an optimal smile is characterized by an upper lip that reaches the gingival margins, with an upward or straight curvature between the philtrum and commissures; an upper incisal line coincident with the border of the lower lip; minimal or no lateral negative space; a commissural line and occlusal frontal plane parallel to the pupillary line; and harmoniously integrated dental and gingival components. However, without intervention, it is infrequent that patients present with all of these components in place. Comprehensive assessment of the deviations from “ideal” and a robust understanding of patient needs and desires allow for development of a targeted, interdisciplinary treatment plan to address patient needs. Patient complaints that are limited to tooth shape and/or shade only can generally be addressed with restorative therapies. Other patient complaints may require further investigation. For example, short clinical tooth crowns may be associated with wear or altered passive eruption. Determination of ideal therapy requires an assessment of incisal display and lip position at rest and in full smile.^{2,70} In cases of little to no incisor display at rest with a

normal lip line during full smile, this can be attributed to limited crown height incisally, and the crowns may be extended using restorative techniques. Conversely, if short clinical crowns are associated with excessive gingival display and normal incisor display at rest, the patient will likely require resective surgical intervention to perform a gingivectomy or a crown-lengthening procedure, depending upon the classification of altered passive eruption. We will review two common patient esthetic complaints and potential interventions below.

Excessive gingival display/“gummy smile”

Excessive gingival display may be due to one or more of the following factors: 1) gingival overgrowth, 2) altered passive eruption, 3) vertical maxillary excess (VME), 4) short upper lip, 5) hypermobile upper lip, and 6) dentoalveolar extrusion.⁷⁰ Clinical examination to determine clinical crown height, lip length and mobility, and vertical facial-third symmetry should allow for identification of the underlying etiology for the gingival display (figure 2).^{71,72} In cases of gingival overgrowth and altered passive eruption, patients will present with short clinical crowns and either excessive gingival tissues or excessive gingiva and alveolar bone, respectively.² Patients with altered passive eruption demonstrate periodontal attachment apparatus at a position that is more coronal than anatomic norms. These patients should be treated with resective surgical intervention and an evaluation of potential underlying systemic conditions or medications that may contribute to gingival overgrowth.⁷³ Both VME and a short upper lip may present with lip incompetence, but lip length will be normal in cases of VME.² VME is associated with a lengthening of the lower facial third. VME may be addressed with orthognathic surgery, and a short upper lip would require lip lengthening surgery to reduce gingival display.⁷² Lip hypermobility is present in cases of normal lip length with increased mobility. Addressing this may require lip repositioning surgery and/or, in some cases, treatment with botulinum toxin to reduce muscle hyperfunction.⁷²⁻⁷⁴ Dentoalveolar extrusion occurs due to overeruption of maxillary incisors and may present with increased clinical crown length, potential lip incompetence, and alteration of arch curvature.⁷² Treatment should include orthodontic intrusion of the supererupted incisors. In some cases, patients may present with concomitant underlying etiologies for excessive gingival display and coordinated interdisciplinary care may be necessary.⁷²

Smile asymmetry

Symmetry has been associated with increased perceived facial attractiveness.^{75,76} Patient histories and complaints regarding symmetry should be carefully reviewed to determine patients’ concern and the rationale for proposed therapy (figure 3). Asymmetry associated with tooth shape may be addressed restoratively or may require orthodontic tooth movement to establish space to allow for ideal restorative care. Asymmetry associated with midline, smile arch, or smile cant may require treatment with orthodontic and/or orthognathic means.⁷⁷ Gingival contour



FIGURE 3: Assessment of symmetry of the teeth related to the facial midline as well as ideal tooth proportions can indicate a need for interdisciplinary care.

asymmetry may be due to gingival recession or gingival overgrowth. Complete documentation of examination findings and thorough assessment of gingival margin relationship with the cemento-enamel junction is critical to appropriate diagnosis.⁷⁸ Root coverage periodontal plastic surgery procedures, either alone or in combination with restorative therapies, can be used to reestablish appropriate gingival contours.⁷⁷ Lip asymmetry may be associated with volume and/or mobility. Asymmetry associated with lip volume may be addressed with dermal fillers,^{79,80} and myofascial exercises may be helpful to strengthen weakened muscles leading to asymmetrical lip movement.⁸¹

CONCLUSION

Understanding the complex interactions of the components of smile esthetics and the potential interventions that can be provided by dental health-care professionals leads to optimal outcomes. Care must be patient-centered to address individual patients' perceived esthetic challenges. Demonstration of likely treatment outcomes using digital modeling and/or reversible procedures may allow patients and practitioners to have a robust discussion about predictable and achievable outcomes of treatment. It is also important to consider interdisciplinary treatment modalities that provide solutions to address overlapping etiologies that contribute to compromised smile esthetics.

REFERENCES

1. American Academy of Orthodontists Press Releases. A good smile can lead to improved love, social and professional success, according to the American Association of Orthodontists. November 11, 2014. <http://couserorthodontics.com/study-shows-that-one-third-of-american-adults-are-unhappy-with-their-smile/>. Accessed June 11, 2020.
2. Garber DA, Salama MA. The aesthetic smile: diagnosis and treatment. *Periodontol* 2000. 1996;11:18-28.
3. Preston JD. The golden proportion revisited. *J Esthet Dent*. 1993;5:247-251.
4. Levin EI. Dental esthetics and the golden proportion. *J Prosthet Dent*. 1978; 40(3):244-252.
5. Qualtrough AJ, Burke FJ. A look at dental esthetics. *Quintessence Int*. 1994;25(1):7-14.
6. Zawawi KH, Malki GA, Al-Zahrani MS, Alkhiari YM. Effect of lip position and gingival display on smile and esthetics as perceived by college students with different educational backgrounds. *Clin Cosmet Investig Dent*. 2013;5:77-80.
7. Afroz S, Rathi S, Rajput G, Rahman SA. Dental esthetics and its impact on psycho-social well-being and dental self-confidence: a campus-based survey of north Indian university students. *J Indian Prosthodont Soc*. 2013;13(4):455-460.
8. Terry DA, Geller W, Tric O, et al. Anatomical form defines color, function, form and aesthetics. *Pract Proced Aesthet Dent*. 2002;14(1):59-67.
9. Hasegawa A, Ikeda I, Kawaguchi S. Color and translucency of in vivo natural central incisors. *J Prosthet Dent*. 2000;83(4):418-423.
10. Esan TA, Olusile AO, Akeredolu PA. Factors influencing tooth shade selection for completely edentulous patients. *J Contemp Dent Pract*. 2006;7(5):80-87.
11. Niaz MO, Naseem M, Elcock C. Ethnicity and perception of dental shade esthetics. *Int J Esthet Dent*. 2015;10(2):286-298.
12. Söderfeldt B, Palmqvist S, Arnbjerg D. Factors affecting attitudes toward dental appearance and dental function in a Swedish population aged 45–69 years. *Community Dent Health*. 1993;10:123-130.
13. Peck S, Peck L, Kataja M. Some vertical lineaments of lip position. *Am J Orthod Dentofacial Orthop*. 1992;101(6):519-524.
14. Gillen RJ, Schwartz RS, Hilton TJ, Evans DB. An analysis of selected normative tooth proportions. *Int J Prosthodont*. 1994;7(5):410-417.
15. German DS, Chu SJ, Furlong ML, Patel A. Simplifying optimal tooth-size calculations and communications between practitioners. *Am J Orthod Dentofacial Orthop*. 2016;150(6):1051-1055.
16. Sarver DM, Ackerman MB. Dynamic smile visualization and quantification: part 2. Smile analysis and treatment strategies. *Am J Orthod Dentofacial Orthop*. 2003;124(2):116-127.
17. Lombardi RE. The principles of visual perception and their clinical application to denture esthetics. *J Prosth Dent*. 1973;29(4):358-382.
18. Kokich VO Jr, Kiyak HA, Shapiro PA. Comparing perceptions of dental and lay people to altered dental esthetics. *J Esthet Dent*. 1999;11(6):311-324.
19. Sarver DM. The importance of incisor positioning in the esthetic smile: the smile arc. *Am J Orthod Dentofacial Orthop*. 2001;120(2):98-111.
20. Frush JP, Fisher RD. The dynesthetic interpretation of the dentogenic concept. *J Prosth Dent*. 1958;8(4):558-581.
21. Dong JK, Jin TH, Cho HW, Oh SC. The esthetics of the smile: a review of some recent studies. *Int J Prosthodont*. 1999;12(1):9-19.
22. Rigsbee OH 3rd, Sperry TP, BeGole EA. The influence of facial animation on smile characteristics. *Int J Adult Orthodon Orthognath Surg*. 1988;3(4):233-239.
23. Ackerman MB, Ackerman JL. Smile analysis and design in the digital era. *J Clin Orthod*. 2002;36(4):221-236.
24. Zachrisson BU. Esthetic factors involved in anterior tooth display and the smile: vertical dimension. *J Clin Orthod*. 1998;32:432-445.
25. Philips E. The classification of smile patterns. *J Can Dent Assoc*. 1999;65:252-254.
26. Hulseley CM. An esthetic evaluation of lip-teeth relationships present in the smile. *Am J Orthod*. 1970;57(2):32-44.
27. Machado AW. 10 commandments of smile esthetics. *Dent Press J Orthod*. 2014;19(4):136-157.

28. Machado AW, Santos TC, Araujo TM, Gandini LG. Integração Ortodontia e Dermatologia na busca da excelência na estética labial. *Rev Clin Ortodon Dent Press*. 2010;9(3):47-56.
29. American Society of Plastic Surgeons. 2017 cosmetic plastic surgery statistics. <https://www.plasticsurgery.org/documents/News/Statistics/2017/plastic-surgery-statistics-report-2017.pdf>. Accessed August 4, 2020.
30. Orban B. Histology and physiology of the gingiva. *J Am Dent Assoc*. 1952;44(6):624-628.
31. Morley J, Eubank J. Macroesthetic elements of smile design. *J Am Dent Assoc*. 2001;132(1):39-45.
32. Tarnow DP, Magner AW, Fletcher P. The effect of the distance from the contact point to the crest of bone on the presence or absence of the interproximal dental papilla. *J Periodontol*. 1993;63(12):995-996.
33. Sabri R. Treatment of a class I malocclusion with an ankylosed maxillary central incisor. *Am J Orthod Dentofacial Orthop*. 2002;122(5):557-565.
34. Dolt AH 3rd, Robbins JW. Altered passive eruption: an etiology of short clinical crowns. *Quintessence*. 1997;28(6):363-372.
35. Løe H, Ånerud Å, Boysen H. The natural history of periodontal disease in man: prevalence, severity, and extent of gingival recession. *J Periodontol*. 1992;63(6):489-495.
36. Ahmad I. Geometric considerations in anterior dental aesthetics: restorative principles. *Pract Periodontics Aesthet Dent*. 1998;10:813-822.
37. Mackley RJ. An evaluation of smiles before and after orthodontic treatment. *Angle Orthod*. 1993;63(3):183-189.
38. Geron S, Atalia W. Influence of sex on the perception of oral and smile esthetics with different gingival display and incisal plane inclination. *Angle Orthod*. 2005;75(5):778-784.
39. Ackerman MB, Brensinger C, Landis JR. An evaluation of dynamic lip-tooth characteristics during speech and smile in adolescents. *Angle Orthod*. 2004;74(1):43-50.
40. Hunt O, Johnston C, Hepper P, et al. The influence of maxillary gingival exposure on dental attractiveness ratings. *Eur J Orthod*. 2002;24(2):199-204.
41. Keating CF. Gender and the physiognomy of dominance and attractiveness. *Soc Psychol Quart*. 1985;48(1):61-70.
42. Bashour M. History and current concepts in the analysis of facial attractiveness. *Plast Reconstr Surg*. 2006;118(3):741-756.
43. Cunningham MR. Measuring the physical in physical attractiveness: quasi-experiments on the sociobiology of female facial beauty. *J Personal Soc Psychol*. 1986;50(5):925-935.
44. Cunningham MR, Barbee AP, Pike CL. What do women want? Facialmetric assessment of multiple motives in the perception of male facial attractiveness. *J Personal Soc Psychol*. 1990;59(1):61-72.
45. Riggio RE, Widaman KF, Tucker JS, Salinas C. Beauty is more than skin deep: components of attractiveness. *Bas Appl Soc Psychol*. 1991;12(4):423-439.
46. Shaw WC. The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am J Orthod*. 1981;79(4):399-415.
47. Foo P, Sampson W, Roberts R, et al. Facial aesthetics and perceived need for further treatment among adults with repaired cleft as assessed by cleft team professionals and laypersons. *Eur J Orthod*. 2013;35(3):341-346.
48. Rhodes G, Sumich A, Byatt G. Are average facial configurations attractive only because of their symmetry? *Psychol Sci*. 1999;10(1):52-58.
49. Mealey L, Bridgstock R, Townsend GC. Symmetry and perceived facial attractiveness: a monozygotic co-twin comparison. *J Personal Soc Psychol*. 1999;76(1):151-158.
50. Hönn M, Göz G. The ideal of facial beauty: a review. *J Orofac Orthop*. 2007;68(1):6-16.
51. Alsulaimani FF, Batwa W. Incisors' proportions in smile esthetics. *J Orthod Sci*. 2013;2(3):109-112.
52. Rubin LR. The anatomy of a smile: its importance in the treatment of facial paralysis. *Plast Reconstr Surg*. 1974;53(4):384-387.
53. Paletz JL, Manktelow RT, Chaban R. The shape of a normal smile: implications for facial paralysis reconstruction. *Plast Reconstr Surg*. 1994;93(4):784-789.
54. Sarver DM, Ackerman MB. Dynamic smile visualization and quantification: part 1. Evolution of the concept and dynamic records for smile capture. *Am J Orthod Dentofacial Orthop*. 2003;124(1):4-12.
55. Miller CJ. The smile line as a guide to anterior esthetics. *Dent Clin N Am*. 1989;33:157-164.
56. Ackerman JL, Ackerman MB, Brensinger CM, Landis JR. A morphometric analysis of the posed smile. *Clin Orthod Res*. 1998;1(1):2-11.
57. Gan J, Li L, Zhai Y, Liu Y. Deep self-taught learning for facial beauty prediction. *Neurocomputing*. 2014;144:295-303.
58. Xie D, Liang L, Jin L, et al. SCUT-FBP: a benchmark dataset for facial beauty perception. In: *2015 IEEE international conference on systems, man, and cybernetics IEEE*. 2015:1821-1826.
59. Ras F, Habets LL, van Ginkel FC, Prah Andersen B. Quantification of facial morphology using stereophotogrammetry—demonstration of a new concept. *J Dent*. 1996;24(5):369-374.
60. Proffit WR, Sarver DM, Ackerman JL. Diagnosis and treatment planning. In: Proffit WR, Fields HW, Sarver DM, eds. *Contemporary Orthodontics*. 4th ed. Mosby Elsevier; 2007:167-233.
61. Kiyak HA. Does orthodontic treatment affect patients' quality of life? *J Dent Educ*. 2008;72(8):886-894.
62. Flores-Mir C, Silva E, Barriga MI, et al. Lay person's perception of smile aesthetics in dental and facial views. *J Orthod*. 2004;31(3):204-209.
63. McLeod C, Fields HW, Hechter F, et al. Esthetics and smile characteristics evaluated by laypersons. *Angle Orthod*. 2011;81(2):198-205.
64. Ioi H, Kang S, Shimomura T, et al. Effects of vertical positions of anterior teeth on smile esthetics in Japanese and Korean orthodontists and orthodontic patients. *J Esthet Restor Dent*. 2013;25(4):274-282.
65. Ker AJ, Chan R, Fields HW, et al. Esthetics and smile characteristics from the layperson's perspective: a computer-based survey study. *J Am Dent Assoc*. 2008;139(10):1318-1327.
66. Tüzgıray YB, Kaya B. Factors affecting smile esthetics. *Turk J Orthod*. 2013;26(1):58-64.
67. Sriphadungporn C, Chamnannidiadha N. Perception of smile esthetics by laypeople of different ages. *Prog Orthod*. 2017;18(1):8.
68. Russello S. The impact of media exposure on self-esteem and body satisfaction in men and women. *J Interdisciplinary Undergrad Res*. 2009;1(1):1-12.
69. Patnaik VVG, Singla Rajan K, Bala S. Anatomy of 'a beautiful face & smile.' *J Anat Soc India*. 2003;52:74-80.
70. Kokich VG. Esthetics: the orthodontic-periodontic restorative connection. *Seminars Orthod*. 1996;2:21-30.
71. Silberberg N, Goldstein M, Smidt A. Excessive gingival display—etiology, diagnosis, and treatment modalities. *Quintessence Int*. 2009;40(10):809-818.
72. Dym H, Pierre R 2nd. Diagnosis and treatment approaches to a "gummy smile." *Dent Clin North Am*. 2020;64(2):341-349.
73. Abou-Arraj RV, Souccar NM. Periodontal treatment of excessive gingival display. *Seminars Orthod*. 2013;19(4):267-278.
74. Dinker S, Anitha A, Sorake A, Kumar K. Management of gummy smile with botulinum toxin type-A: a case report. *J Int Oral Health*. 2014;6(1):111-115.
75. Penton-Voak IS, Jones BC, Little AC, et al. Symmetry, sexual dimorphism in facial proportions and male facial attractiveness. *Proc Biol Sci*. 2001;268(1476):1617-1623.
76. Grammer K, Thornhill R. Human (Homo sapiens) facial attractiveness and sexual selection: the role of symmetry and averageness. *J Comp Psychol*. 1994;108(3):233-242.
77. Thiesen G, Gribel BF, Freitas MPM. Facial asymmetry: a current review. *Dental Press J Orthod*. 2015;20(6):110-125.
78. Jati AS, Furquim LZ, Consolaro A. Gingival

recession: its causes and types, and the importance of orthodontic treatment. *Dental Press J Orthod.* 2016;21(3):18-29.

79. Vedamurthy M, Vedamurthy A. Dermal fillers: tips to achieve successful outcomes. *J Cutan Aesthet Surg.* 2008;1(2):64-67.
80. Dastoor SF, Misch CE, Wang H-L. Dermal fillers for facial soft tissue augmentation. *J Oral Implantol.* 2007;33(4):191-204.
81. Brach JS, VanSwearingen JM. Physical therapy for facial paralysis: a tailored treatment approach. *Phys Ther.* 1999;79(4):397-404.



MARIA L. GEISINGER, DDS, MS, is a professor and director of advanced education in periodontology in the Department of Periodontology in the University of Alabama at Birmingham (UAB) School of Dentistry. Dr. Geisinger received her BS in biology from Duke University, her DDS from Columbia University School of Dental Medicine, and her MS and certificate in periodontology and implantology from the University of Texas Health Science Center at San

Antonio. Dr. Geisinger is a diplomate in the American Board of Periodontology. She has served as the president of the American Academy of Periodontology Foundation and on multiple national and regional organized dentistry



HUSSEIN BASMA, DDS, DESS, MS, graduated from Damascus University in 2009 with a doctor of dental surgery, after receiving a full scholarship. Following dental school, Dr. Basma completed a graduate degree in prosthodontics at Lebanese University. He completed his certificate in periodontology and his master of science at the University of Alabama at Birmingham (UAB). His research interests focus on bone grafting and implant dentistry. Dr. Basma is

currently a full-time assistant professor in the Department of Periodontology at the UAB School of Dentistry. His clinical and teaching efforts focus on interdisciplinary care in dentistry and the integration of technology to deliver optimal patient care.

QUICK ACCESS CODE 21002

ONLINE COMPLETION

Take this test online for immediate credit. Visit dentalacademyofce.com and sign in. If you have not previously purchased the course, select it from the "Online Courses" listings and complete your purchase. The exam will then be added to your "Archives" page, where a "Take Exam" link will be provided. Click on this link, complete all questions, and submit your answers. An immediate grade report will be generated. If you receive a score of 70% or higher, your verification form will be provided immediately for viewing and printing. View and print forms at any time by visiting the site and returning to your "Archives."

QUESTIONS

- A study commissioned by the American Association of Orthodontists suggests that at least ___ US adults are unhappy with their smile esthetics.**
 - A. 1:2
 - B. 1:3
 - C. 1:5
 - D. 1:10
- Approximately ___ of US adults ages 18-24 report that they have untagged pictures of themselves on social media because they didn't like their smiles.**
 - A. 24%
 - B. 36%
 - C. 48%
 - D. 56%
- The anatomical features of an esthetic smile include all of the following except:**
 - A. Teeth
 - B. Lips
 - C. Gingivae
 - D. Tongue
- In individuals who were unsatisfied with their smile esthetics, the most common complaint was:**
 - A. Tooth size
 - B. Tooth position
 - C. Tooth color
 - D. Lip shape
- The average vertical height of maxillary central incisors is ___ in males and females, respectively.**
 - A. 10.6 mm and 9.8 mm
 - B. 11.2 mm and 10.6 mm
 - C. 9.8 mm and 8.7 mm
 - D. 12.3 mm and 11.2 mm
- The ideal width-to-length ratio for teeth is considered to be the "golden proportion." The mean width-to-length ratio for teeth in adults in a range of ___ is judged as esthetically acceptable.**
 - A. 80%-85%
 - B. 75%-80%
 - C. 70%-75%
 - D. 65%-70%
- The dental midline at the maxillary central incisors that corresponds to the facial midline (a line from the nasion to the base of the philtrum, or "cupid's bow") is deemed most esthetic.**
 - A. Both statements are true.
 - B. The first statement is true; the second statement is false.
 - C. The first statement is false; the second statement is true.
 - D. Both statements are false.
- Mild midline discrepancies in cases with other esthetic smile components and a vertically oriented interproximal contact between the maxillary central incisors are judged unacceptable by dentists and laypeople.**
 - A. Both statements are true.
 - B. The first statement is true; the second statement is false.
 - C. The first statement is false; the second statement is true.
 - D. Both statements are false.

ONLINE COMPLETION

Take this test online for immediate credit. Visit dentalacademyofce.com and sign in. If you have not previously purchased the course, select it from the "Online Courses" listings and complete your purchase. The exam will then be added to your "Archives" page, where a "Take Exam" link will be provided. Click on this link, complete all questions, and submit your answers. An immediate grade report will be generated. If you receive a score of 70% or higher, your verification form will be provided immediately for viewing and printing. View and print forms at any time by visiting the site and returning to your "Archives."

QUESTIONS

8. **A. Lack of buccal corridor fill has been determined to be detectable by laypersons as a drive of esthetics. B. Buccal corridor fill may also be influenced by the relative antero-posterior position of the maxilla in relation to the frame of the lips.**
- A. Both statements are true.
 B. The first statement is true; the second statement is false.
 C. The first statement is false; the second statement is true.
 D. Both statements are false.
9. **Lip length is measured in repose from ___ at the midline and should be approximately equal in length with commissure height.**
- A. The nasion to the most inferior portion of the upper lip
 B. The subnasale point to the philtrum
 C. The nasion to the philtrum
 D. The subnasale point to the most inferior portion of the upper lip
10. **On average, lip length is ___ in males and females, respectively, and has been shown to increase with age.**
- A. 18 mm and 16 mm
 B. 20 mm and 18 mm
 C. 23 mm and 20 mm
 D. 25 mm and 23 mm
11. **In most patients, lip mobility accounts for approximately ___ of the original lip length.**
- A. 60%
 B. 70%
 C. 80%
 D. 90%
12. **While an average 10 mm of maxillary incisal display is present at full smile, the interindividual variation includes a physiologic range from ___.**
- A. 5-15 mm C. 4-10 mm
 B. 2-12 mm D. 3-14 mm
13. **Papillary fill can be predicted by the relationship of interproximal bone levels to the apical extent of the interdental contact point. If the distance from the interdental contact point to the crest of the interproximal bone is £5 mm, papillae are present 100% of the time, but the prevalence of papillae when the distance between the contact point and the interproximal crest is 6 mm is ___.**
- A. 72% C. 56%
 B. 64% D. 47%
14. **The height of gingival contour of the central incisors generally corresponds to that of the canines with the gingival margins of the lateral incisors approximately ___ more coronal than those of the canines/central incisors.**
- A. 3.5-4 mm C. 2-2.5 mm
 B. 2.5-3 mm D. 1-1.5 mm
15. **A. While the ideal amount of gingival display is not uniform among populations, some investigators have shown that ideal lip position occurs when the lower margin of the upper lip aligns evenly with the gingival margin of the maxillary central incisors. B. Gingival display is judged to be more detrimental to esthetics in men when compared to women.**
- A. Both statements are true.
 B. The first statement is true; the second statement is false.
 C. The first statement is false; the second statement is true.
 D. Both statements are false.
16. **All of the following are associated with facial attractiveness, except:**
- A. Quality of dating partners
 B. Self-reported health outcomes
 C. Perceived intelligence
 D. Perceived pleasantness
17. **The presence of a unilateral lateral incisor width change of > ___ is considered unesthetic by both dentists and laypersons.**
- A. 2.5 mm
 B. 1.5 mm
 C. 1 mm
 D. 0.5 mm
18. **A consonant smile arc is defined as:**
- A. Alignment of the maxillary incisal edges with the inner contour of the lower lip at full smile
 B. Alignment of the maxillary incisal edges 2 mm above the lower lip contour
 C. A smile with completely filled buccal corridor space
 D. Alignment of gingival zeniths with the inferior border of the upper lip at full smile
19. **Incisal edge and lip curvature tend to be more pronounced in:**
- A. Younger patients
 B. Patients who were treated with orthodontic therapy
 C. Older individuals
 D. A and B
20. **In order to critically assess the individual smile components and their interaction, careful examination and documentation are necessary in both ___ and ___ function.**
- A. Rest and masticatory
 B. Static and dynamic
 C. Masticatory and phonetic
 D. Phonetic and static

ONLINE COMPLETION

Take this test online for immediate credit. Visit dentalacademyofce.com and sign in. If you have not previously purchased the course, select it from the "Online Courses" listings and complete your purchase. The exam will then be added to your "Archives" page, where a "Take Exam" link will be provided. Click on this link, complete all questions, and submit your answers. An immediate grade report will be generated. If you receive a score of 70% or higher, your verification form will be provided immediately for viewing and printing. View and print forms at any time by visiting the site and returning to your "Archives."

QUESTIONS

- 21. A. Age, gender, geographic divergence, and cultural norms may impact perception of smile esthetics. B. Women and younger individuals are more tolerant of certain deviations from esthetic standards.**
- Both statements are true.
 - The first statement is true; the second statement is false.
 - The first statement is false; the second statement is true.
 - Both statements are false.
- 22. Dental health-care professionals and plastic surgeons have a ___ standard for facial esthetics than laypersons with similar demographics and socioeconomic status.**
- Lower
 - Higher
 - Similar
 - More complex
- 23. Patients' individual standards and self-perception are ___ to their smile esthetics when compared to the opinions of their dental health-care providers.**
- Equally as important
 - Less important
 - More important
 - Irrelevant
- 24. A. Over time, esthetic ideals are mutable and may be influenced by mass media, including TV, films, magazines, fashion, advertisements, and social media. B. The esoteric nature of "ideal" attractiveness also increases the importance of initial trials with reversible interventions, including digital modeling, to assess individual patient satisfaction with likely outcomes of therapy and to manage patient expectations for predictable results.**
- Both statements are true.
 - The first statement is true; the second statement is false.
 - The first statement is false; the second statement is true.
 - Both statements are false.
- 25. Which of the following are potential etiologies associated with excessive gingival display ("gummy smile")?**
- Altered passive eruption
 - Vertical maxillary excess (VME)
 - Short upper lip
 - All of the above
- 26. Altered passive eruption is characterized by excessive gingival display, short clinical crowns, and ___ position of the periodontal attachment apparatus.**
- Apical
 - Buccal
 - Coronal
 - Posterior
- 27. Lip incompetence at repose is noted in all of the following conditions except:**
- Vertical maxillary excess (VME)
 - Gingival overgrowth
 - Short maxillary lip
 - Dentoalveolar extrusion
- 28. Vertical maxillary excess is associated with which of the following findings?**
- Normal maxillary lip length
 - Elongated lower facial third
 - Lip incompetence at repose
 - All of the above
- 29. Gingival contour asymmetry due to gingival recession can be identified through careful assessment of gingival margin relationship with the cemento-enamel junction. Root coverage periodontal plastic surgery procedures, either alone or in combination with restorative therapies, can be used to reestablish appropriate gingival contours.**
- Both statements are true.
 - The first statement is true; the second statement is false.
 - The first statement is false; the second statement is true.
 - Both statements are false.
- 30. Demonstration of likely treatment outcomes for patients using ___ can allow patients and practitioners to have a robust discussion about predictable and achievable outcomes of treatment and may direct practitioners to more ideal therapies for individual patients.**
- Reversible procedures
 - Digital modeling
 - Interdisciplinary therapy
 - All of the above

PUBLICATION DATE:	MAY 2021
EXPIRATION DATE:	APRIL 2024

Is beauty truly in the eye of the beholder?

Evidence-based ideal smile esthetics

Name: _____ Title: _____ Specialty: _____

Address: _____ Email: _____ AGD member ID (if applies): _____

City: _____ State: _____ ZIP: _____ Country: _____

Telephone: Primary () _____ Office () _____

Requirements for obtaining CE credits by mail/fax: 1) Read entire course. 2) Complete info above. 3) Complete test by marking one answer per question. 4) Complete course evaluation. 5) Complete credit card info or write check payable to Endeavor Business Media. 6) Mail/fax this page to DACE. A score of 70% is required for CE credit. **For questions, call (800) 633-1681. Course may also be completed at dentalacademyofce.com.**

EDUCATIONAL OBJECTIVES

- List the components of a smile and discuss their roles in optimal smile esthetics
- Describe the step-by-step approach to evaluating a patient's smile, including assessing tooth shape/shade, gingival display, and lip length/mobility
- Understand the role of patient-based characteristics on acceptable esthetics and ideal smile components
- Discuss personalized treatment options to achieve ideal smile esthetics based upon underlying diagnoses in each patient

COURSE EVALUATION

1. Were the individual course objectives met?

Objective #1: Yes No Objective #2: Yes No

Objective #3: Yes No Objective #4: Yes No

Please evaluate this course by responding to the following statements, using a scale of Excellent = 5 to Poor = 0.

2. To what extent were the course objectives accomplished overall?	5	4	3	2	1	0
3. Please rate your personal mastery of the course objectives.	5	4	3	2	1	0
4. How would you rate the objectives and educational methods?	5	4	3	2	1	0
5. How do you rate the author's grasp of the topic?	5	4	3	2	1	0
6. Please rate the instructor's effectiveness.	5	4	3	2	1	0
7. Was the overall administration of the course effective?	5	4	3	2	1	0
8. Please rate the usefulness and clinical applicability of this course.	5	4	3	2	1	0
9. Please rate the usefulness of the supplemental bibliography.	5	4	3	2	1	0
10. Do you feel that the references were adequate?		Yes		No		
11. Would you participate in a similar program on a different topic?		Yes		No		

12. If any of the continuing education questions were unclear or ambiguous, please list them.

13. Was there any subject matter you found confusing? Please describe.

14. How long did it take you to complete this course?

15. What additional continuing dental education topics would you like to see?

Mail/fax completed answer sheet to:

Endeavor Business Media

Attn: Dental division

7666 E. 61st St. Suite 230, Tulsa, OK 74133

Fax: (918) 831-9804

Payment of \$59 is enclosed.

Make check payable to Endeavor Business Media

If paying by credit card, please complete the following: MC Visa AmEx Discover

Acct. number: _____

Exp. date: _____ CVC #: _____

Billing address: _____

Charges on your statement will show up as PennWell / Endeavor.

1. (A) (B) (C) (D)	16. (A) (B) (C) (D)
2. (A) (B) (C) (D)	17. (A) (B) (C) (D)
3. (A) (B) (C) (D)	18. (A) (B) (C) (D)
4. (A) (B) (C) (D)	19. (A) (B) (C) (D)
5. (A) (B) (C) (D)	20. (A) (B) (C) (D)
6. (A) (B) (C) (D)	21. (A) (B) (C) (D)
7. (A) (B) (C) (D)	22. (A) (B) (C) (D)
8. (A) (B) (C) (D)	23. (A) (B) (C) (D)
9. (A) (B) (C) (D)	24. (A) (B) (C) (D)
10. (A) (B) (C) (D)	25. (A) (B) (C) (D)
11. (A) (B) (C) (D)	26. (A) (B) (C) (D)
12. (A) (B) (C) (D)	27. (A) (B) (C) (D)
13. (A) (B) (C) (D)	28. (A) (B) (C) (D)
14. (A) (B) (C) (D)	29. (A) (B) (C) (D)
15. (A) (B) (C) (D)	30. (A) (B) (C) (D)

PLEASE PHOTOCOPY ANSWER SHEET FOR ADDITIONAL PARTICIPANTS.

INSTRUCTIONS

All questions have only one answer. If mailed or faxed, grading of this examination is done manually. Participants will receive confirmation of passing by receipt of a Verification of Participation form. The form will be mailed within two weeks after receipt of an examination.

COURSE EVALUATION AND FEEDBACK

We encourage participant feedback. Complete the evaluation above and e-mail additional feedback to Aileen Southerland (asoutherland@endeavor2b.com) and Laura Winfield (lwinfield@endeavor2b.com).

COURSE CREDITS AND COST

All participants scoring 70% or higher on the examination will receive a verification form for three (3) continuing education (CE) credits. Participants are urged to contact their state dental boards for CE requirements. The cost for courses ranges from \$20 to \$110.

PROVIDER INFORMATION

Endeavor Business Media is an ADA CERP-recognized provider. ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of continuing dental education. ADA CERP neither approves nor endorses individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry. Concerns about a CE provider may be directed to the provider or to ADA CERP at ada.org/gotocepr.

Endeavor Business Media is designated as an approved PACE program provider by the Academy of General Dentistry. The formal continuing dental education programs of this program provider are accepted by the AGD for fellowship, mastership, and membership maintenance credit. Approval does not imply acceptance by a state or provincial board of dentistry or AGD endorsement. The current term of approval extends from 11/1/2019 to 10/31/2022. Provider ID# 320452. AGD code: 780.

Dental Board of California: Provider RP9933. Course registration number CA code: 03-5933-21002. Expires 7/31/2022. This course meets the Dental Board of California's requirements for three (3) units of continuing education.

Endeavor Business Media is designated as an approved provider by the American Academy of Dental Hygiene Inc. #AADHPNW (January 1, 2020 - December 31, 2021). Approval does not imply acceptance by a state or provincial board of dentistry. Licensee should maintain this document in the event of an audit.

RECORD KEEPING

Endeavor Business Media maintains records of your successful completion of any exam for a minimum of six years. Please contact our offices for a copy of your CE credits report. This report, which will list all credits earned to date, will be generated and mailed to you within five business days of receipt.

CANCELLATION AND REFUND POLICY

Participants who are not 100% satisfied can request a refund by contacting Endeavor Business Media in writing.

IMAGE AUTHENTICITY

The images in this educational activity have not been altered.

© 2020 Academy of Dental Therapeutics and Stomatology, a division of Endeavor Business Media